

GastroDoxs
defenders of the digestive system

Intestinal Lymphangiectasia

A Practical Guide For Patients

832-632-4070
info@gastrodoxs.com

TABLE OF CONTENTS

01 Introduction

02 What Causes Intestinal Lymphangiectasia?

03 Symptoms

04 Diagnosis

05 Diet

06 Medications

07 Therapies

08 Coping

09 FAQs

10 Take Action

Introduction: Intestinal Lymphangiectasia

Intestinal lymphangiectasia is a rare condition in which lymphatic vessels within the small intestine become enlarged. These vessels normally transport lymph, dietary fats, proteins, and immune cells. When they become overly dilated, they may rupture or leak protein-rich lymph into the intestine. This leakage can cause protein-losing enteropathy, resulting in low blood protein, swelling, diarrhea, nutritional deficiencies, and reduced immune-cell levels



Key Takeaways

- Intestinal lymphangiectasia involves enlarged lymphatic vessels in the small intestine.
- The enlarged vessels may leak lymph into the digestive tract.
- Protein loss can cause low albumin and widespread swelling.
- Primary intestinal lymphangiectasia is also known as Waldmann disease.
- Secondary intestinal lymphangiectasia results from another medical condition.
- Swelling of the legs is one of the most common signs.
- Diarrhea, abdominal pain, fatigue, and poor weight gain may occur.

What Causes Intestinal Lymphangiectasia?

Known Causes

Intestinal lymphangiectasia develops when lymph cannot flow normally through the intestinal lymphatic system. The pressure causes small lymphatic channels, called lacteals, to expand and leak their contents into the intestine. Primary disease results from abnormal lymphatic development. Secondary disease develops when another condition obstructs lymph flow or increases pressure within the lymphatic circulation.



Common Signs & Symptoms

Symptoms vary according to how much protein and lymph are lost. Some patients have mild disease with few symptoms, while others develop significant swelling, diarrhea, nutritional deficiencies, or fluid accumulation.

Core Symptoms Include:



- Swelling of the feet, ankles, or lower legs
- Generalized body swelling
- Mild or persistent diarrhea
- Abdominal pain or discomfort
- Abdominal bloating
- Nausea
- Fatigue or reduced energy
- Poor appetite

Severe cases can also cause:



- Severe generalized swelling, called anasarca
- Fluid accumulation in the abdomen
- Chylous ascites containing lymphatic fluid
- Fluid around the lungs
- Fluid around the heart
- Shortness of breath
- Severe malnutrition
- Muscle wasting

GastroDoxs
defenders of the digestive system

Diagnosis & Medical Testing

Diagnosis involves confirming protein loss through the intestine, identifying enlarged intestinal lymphatic vessels, and determining whether the condition is primary or secondary. Other causes of low albumin, such as liver disease, kidney disease, severe malnutrition, and inflammatory intestinal disease, must also be evaluated.

Diagnostic Tools

- **Medical history evaluation:** Reviews swelling, diarrhea, growth, weight changes, infections, heart disease, and previous surgery.
- **Physical examination:** Checks leg swelling, abdominal fluid, muscle loss, skin changes, and nutritional status.
- **Complete blood count:** Evaluates anemia, infection, and lymphocyte levels.
- **Serum albumin and total protein:** Detect protein depletion caused by intestinal loss.
- **Immunoglobulin testing:** Identifies low antibody levels caused by lymph leakage.
- **Electrolyte and calcium testing:** Checks for mineral abnormalities.
- **Vitamin testing:** Evaluates vitamins A, D, E, K, B12, folate, and other deficiencies.



Diet

Dietary treatment is the cornerstone of care. Long-chain dietary fats enter intestinal lymphatic vessels and can increase pressure within dilated lacteals. A low-fat diet reduces lymphatic engorgement and leakage. Medium-chain triglycerides are absorbed more directly into the portal bloodstream and place less demand on intestinal lymphatic vessels.

Key Insights

- Follow a strict low-fat diet when recommended.
- Limit long-chain dietary fats that enter the intestinal lymphatic system.
- Use prescribed medium-chain triglyceride oil or MCT-containing products.
- Introduce MCT oil gradually to reduce cramping or diarrhea.
- Eat a high-protein diet to replace intestinal protein loss.
- Include lean poultry, fish, eggs, low-fat dairy, beans, tofu, or other tolerated proteins.
- Eat smaller, frequent meals when appetite is reduced.
- Maintain adequate calories to prevent weight and muscle loss.
- Drink sufficient fluids according to the treatment plan.
- Replace electrolytes during persistent diarrhea.







Medications & Treatment Options

There is no single medicine that corrects every form of intestinal lymphangiectasia. Dietary therapy remains the primary treatment. Medications and infusions may be considered when diet alone does not adequately control protein loss, swelling, or complications. Secondary disease requires treatment of the heart, inflammatory, infectious, malignant, or obstructive condition responsible for abnormal lymph flow.

Treatment Options

- **Low-fat nutritional therapy:** Reduces pressure and leakage from intestinal lymphatic vessels.
- **Medium-chain triglyceride supplementation:** Provides absorbable calories while bypassing much of the intestinal lymphatic pathway.
- **High-protein nutrition:** Replaces protein lost through the digestive tract.
- **Albumin infusion:** Temporarily increases blood albumin in patients with severe swelling or fluid accumulation.
- **Intravenous fluids:** Treat dehydration and circulation problems.



Advanced Therapies & Procedures

Advanced therapy is considered when dietary treatment is unsuccessful, disease is localized to a short intestinal segment, or abnormal lymphatic channels can be treated directly. Imaging the lymphatic system helps determine whether disease is focal, diffuse, or connected to abnormal lymphatic vessels elsewhere in the body.

Therapy Options

- **Magnetic resonance lymphangiography:** Identifies abnormal lymphatic pathways and the extent of disease.
- **Intranodal lymphangiography:** Uses contrast material to map lymphatic leakage or abnormal connections.
- **Lymphatic embolization:** Blocks selected leaking or abnormal lymphatic vessels.
- **Segmental intestinal resection:** Removes a short, severely affected intestinal segment.
- **Repeat upper endoscopy:** Monitors mucosal findings and obtains follow-up biopsies.
- **Capsule endoscopy:** Evaluates the extent of small-intestinal involvement.
- **Double-balloon enteroscopy:** Examines and biopsies areas beyond the reach of routine endoscopy.
- **Drainage of ascites:** Removes severe abdominal fluid when medically necessary.
- **Pleural drainage:** Treats a large fluid collection around the lungs.
- **Pericardial drainage:** Treats significant fluid around the heart.
- **Enteral nutrition:** Provides specially formulated nutrition through the digestive tract.

Coping With Intestinal Lymphangiectasia

Living with intestinal lymphangiectasia may require long-term dietary restriction, supplements, repeated blood testing, and monitoring for swelling or nutritional deficiencies. Symptoms may improve substantially when the diet is followed consistently, but protein loss can return if treatment is interrupted.

Practical Strategies

- Follow the low-fat, high-protein diet consistently.
- Use MCT supplements exactly as directed.
- Work with a registered dietitian familiar with protein-losing enteropathy.
- Monitor body weight regularly.
- Record swelling of the feet, legs, hands, or abdomen.
- Track diarrhea, abdominal pain, and food tolerance.
- Take prescribed vitamins and minerals consistently.
- Complete regular albumin, protein, lymphocyte, and vitamin testing.
- Monitor growth carefully in children and adolescents.

FAQS About Intestinal Lymphangiectasia



Q1: What is intestinal lymphangiectasia?

A: Intestinal lymphangiectasia is a condition in which lymphatic vessels in the small intestine become enlarged and leak protein-rich lymph into the bowel. This can lead to protein-losing enteropathy, low albumin, swelling, diarrhea, and nutritional deficiencies.

Q2: What causes intestinal lymphangiectasia?

A: Primary intestinal lymphangiectasia results from abnormal lymphatic development. Secondary intestinal lymphangiectasia may result from heart disease, lymphoma, Crohn's disease, tuberculosis, sarcoidosis, prior Fontan surgery, or another condition that blocks lymph flow or increases lymphatic pressure.

Q3: How is intestinal lymphangiectasia diagnosed?

A: Diagnosis may involve blood tests showing low albumin and lymphocytes, stool alpha-1 antitrypsin testing to confirm protein loss, upper endoscopy with small-intestinal biopsy, capsule endoscopy, enteroscopy, and lymphatic imaging. Testing is also needed to exclude kidney, liver, heart, inflammatory, infectious, and malignant causes.

Q4: How is intestinal lymphangiectasia treated?

A: The main treatment is a low-fat, high-protein diet supplemented with medium-chain triglycerides. Vitamins, minerals, albumin infusions, octreotide, or specialist-directed medicines may also be needed.

Take Action

If you experience persistent swelling of the legs, unexplained low albumin, chronic diarrhea, abdominal swelling, poor growth, weight loss, or recurring nutritional deficiencies, do not ignore the symptoms. Intestinal lymphangiectasia requires careful evaluation to confirm protein loss and determine whether the condition is primary or caused by another disease.

Visit **GastroDoxs.com** to schedule an appointment and access more resources on digestive health.



Your health matters
take the next step
today!

SCHEDULE AN APPOINTMENT

832-632-4070

info@gastrodoxs.com

GastroDoxs
defenders of the digestive system